

Exotic nilpotent cones and Springer representations of Weyl groups

Anthony Henderson

University of Sydney

June 2010

Outline

The Springer correspondence

Kato's exotic Springer correspondence

Geometry of the exotic nilpotent cone

The Springer correspondence

We consider algebraic groups over \mathbb{C} . The nilpotent cone of GL_n is

$$\mathcal{N}(\mathfrak{gl}_n) = \{x \in \text{Mat}_n \mid x \text{ nilpotent, i.e. all eigenvalues } 0\}.$$

GL_n acts by conjugation: $g.x = gxg^{-1}$.

The Springer correspondence

We consider algebraic groups over \mathbb{C} . The nilpotent cone of GL_n is

$$\mathcal{N}(\mathfrak{gl}_n) = \{x \in \text{Mat}_n \mid x \text{ nilpotent, i.e. all eigenvalues } 0\}.$$

GL_n acts by conjugation: $g.x = gxg^{-1}$.

Theorem (Jordan canonical form)

The GL_n -orbits in $\mathcal{N}(\mathfrak{gl}_n)$ are in bijection with \mathcal{P}_n , the set of partitions of n . For $\lambda \in \mathcal{P}_n$, the orbit \mathcal{O}_λ consists of those $x \in \mathcal{N}(\mathfrak{gl}_n)$ whose Jordan form has blocks of sizes $\lambda_1, \lambda_2, \dots$.

The Springer correspondence

We consider algebraic groups over \mathbb{C} . The nilpotent cone of GL_n is

$$\mathcal{N}(\mathfrak{gl}_n) = \{x \in \text{Mat}_n \mid x \text{ nilpotent, i.e. all eigenvalues } 0\}.$$

GL_n acts by conjugation: $g.x = gxg^{-1}$.

Theorem (Jordan canonical form)

The GL_n -orbits in $\mathcal{N}(\mathfrak{gl}_n)$ are in bijection with \mathcal{P}_n , the set of partitions of n . For $\lambda \in \mathcal{P}_n$, the orbit \mathcal{O}_λ consists of those $x \in \mathcal{N}(\mathfrak{gl}_n)$ whose Jordan form has blocks of sizes $\lambda_1, \lambda_2, \dots$.

Since \mathcal{P}_n also labels irreducible representations V_λ of S_n , we have

$$GL_n \backslash \mathcal{N}(\mathfrak{gl}_n) \longleftrightarrow \{\text{irreducible representations of } S_n\}.$$

This bijection can be defined using a certain action of S_n on the cohomology of Springer fibres: we have $H^{\text{top}}(\mathcal{B}_x) \cong V_\lambda$ if $x \in \mathcal{O}_\lambda$.

For G a connected reductive group, and B a Borel subgroup, define

$$\pi : G \times_B \mathfrak{b} \rightarrow \mathfrak{g} : (g, x) \mapsto g.x.$$

Then the **Springer fibre** $\mathcal{B}_x = \pi^{-1}(x)$ can be identified with

(in general) $\{\text{Borel subalgebras } \mathfrak{b}' \subset \mathfrak{g} \mid x \in \mathfrak{b}'\}$

($G = GL_n$) $\{0 = V_0 \subset V_1 \subset \cdots \subset V_n = \mathbb{C}^n \mid x(V_i) \subseteq V_i\}$

($G = Sp_{2n}$) $\{0 = V_0 \subset V_1 \subset \cdots \subset V_{2n} = \mathbb{C}^{2n} \mid$
 $\langle V_i, V_{2n-i} \rangle = 0, x(V_i) \subseteq V_i\}.$

For G a connected reductive group, and B a Borel subgroup, define

$$\pi : G \times_B \mathfrak{b} \rightarrow \mathfrak{g} : (g, x) \mapsto g \cdot x.$$

Then the **Springer fibre** $\mathcal{B}_x = \pi^{-1}(x)$ can be identified with

(in general) $\{\text{Borel subalgebras } \mathfrak{b}' \subset \mathfrak{g} \mid x \in \mathfrak{b}'\}$

($G = GL_n$) $\{0 = V_0 \subset V_1 \subset \cdots \subset V_n = \mathbb{C}^n \mid x(V_i) \subseteq V_i\}$

($G = Sp_{2n}$) $\{0 = V_0 \subset V_1 \subset \cdots \subset V_{2n} = \mathbb{C}^{2n} \mid$
 $\langle V_i, V_{2n-i} \rangle = 0, x(V_i) \subseteq V_i\}.$

There are two important restrictions of π :

- ▶ $\pi_{\text{nil}} : G \times_B \mathfrak{n} \rightarrow \mathcal{N}(\mathfrak{g})$ is the **Springer resolution** of the nilpotent cone. (Note that if x is nilpotent, then the condition $x(V_i) \subseteq V_i$ can be rewritten $x(V_i) \subseteq V_{i-1}$.)
- ▶ $\pi_{\text{rs}} : G \times_B \mathfrak{b}_{\text{rs}} \rightarrow \mathfrak{g}_{\text{rs}}$ is a Galois covering of the regular semisimple set with Galois group W , the Weyl group of G . So for $x \in \mathfrak{g}_{\text{rs}}$, W acts simply transitively on \mathcal{B}_x .

Lusztig observed that since π is a small map, the derived push-forward complex $R\pi_*\mathbb{C}$ is the intersection cohomology extension of the local system $(\pi_{rs})_*\mathbb{C}$ on \mathfrak{g}_{rs} . Hence W acts on the complex $R\pi_*\mathbb{C}$, and therefore on $H^i(\mathcal{B}_x)$ for any $x \in \mathfrak{g}$.

Lusztig observed that since π is a small map, the derived push-forward complex $R\pi_*\mathbb{C}$ is the intersection cohomology extension of the local system $(\pi_{rs})_*\mathbb{C}$ on \mathfrak{g}_{rs} . Hence W acts on the complex $R\pi_*\mathbb{C}$, and therefore on $H^i(\mathcal{B}_x)$ for any $x \in \mathfrak{g}$.

Theorem (Borho–MacPherson)

The endomorphism algebra of $R(\pi_{\text{nil}})_\mathbb{C}$ in the derived category on $\mathcal{N}(\mathfrak{g})$ is isomorphic to the group algebra of W . Hence we have a **Springer correspondence** between the irreps of W and the simple constituents of $R(\pi_{\text{nil}})_*\mathbb{C}$, which are intersection cohomology complexes $IC(\overline{\mathcal{O}}, \mathcal{E})$ where \mathcal{O} is a G -orbit in $\mathcal{N}(\mathfrak{g})$ and \mathcal{E} is a G -equivariant simple local system on \mathcal{O} .*

Lusztig observed that since π is a small map, the derived push-forward complex $R\pi_*\mathbb{C}$ is the intersection cohomology extension of the local system $(\pi_{rs})_*\mathbb{C}$ on \mathfrak{g}_{rs} . Hence W acts on the complex $R\pi_*\mathbb{C}$, and therefore on $H^i(\mathcal{B}_x)$ for any $x \in \mathfrak{g}$.

Theorem (Borho–MacPherson)

The endomorphism algebra of $R(\pi_{\text{nil}})_\mathbb{C}$ in the derived category on $\mathcal{N}(\mathfrak{g})$ is isomorphic to the group algebra of W . Hence we have a **Springer correspondence** between the irreps of W and the simple constituents of $R(\pi_{\text{nil}})_*\mathbb{C}$, which are intersection cohomology complexes $IC(\overline{\mathcal{O}}, \mathcal{E})$ where \mathcal{O} is a G -orbit in $\mathcal{N}(\mathfrak{g})$ and \mathcal{E} is a G -equivariant simple local system on \mathcal{O} .*

In general, not all such $IC(\overline{\mathcal{O}}, \mathcal{E})$ occur in $R(\pi_{\text{nil}})_*\mathbb{C}$. But the trivial local systems always do occur, so we get an injective map

$$G \backslash \mathcal{N}(\mathfrak{g}) \hookrightarrow \{\text{irreps of } W\} : G \cdot x \mapsto H^{\text{top}}(\mathcal{B}_x)^{G_x}.$$

The reason this is bijective in the case of GL_n is that all stabilizers are connected, so there are no non-trivial equivariant local systems.

Consider the situation in types B_n and C_n .

Theorem (Gerstenhaber)

Nilpotent orbits of SO_{2n+1} and Sp_{2n} are classified by Jordan form:

$$SO_{2n+1} \setminus \mathcal{N}(\mathfrak{so}_{2n+1}) \leftrightarrow \left\{ \begin{array}{l} \text{partitions of } 2n+1 \text{ in which every} \\ \text{even part occurs with even multiplicity} \end{array} \right\}$$
$$Sp_{2n} \setminus \mathcal{N}(\mathfrak{sp}_{2n}) \leftrightarrow \left\{ \begin{array}{l} \text{partitions of } 2n \text{ in which every} \\ \text{odd part occurs with even multiplicity} \end{array} \right\}$$

Consider the situation in types B_n and C_n .

Theorem (Gerstenhaber)

Nilpotent orbits of SO_{2n+1} and Sp_{2n} are classified by Jordan form:

$$SO_{2n+1} \setminus \mathcal{N}(\mathfrak{so}_{2n+1}) \leftrightarrow \left\{ \begin{array}{l} \text{partitions of } 2n+1 \text{ in which every} \\ \text{even part occurs with even multiplicity} \end{array} \right\}$$
$$Sp_{2n} \setminus \mathcal{N}(\mathfrak{sp}_{2n}) \leftrightarrow \left\{ \begin{array}{l} \text{partitions of } 2n \text{ in which every} \\ \text{odd part occurs with even multiplicity} \end{array} \right\}$$

Example

Take $n = 3$. The possible Jordan types for $\mathcal{N}(\mathfrak{so}_7)$ are:

$$(7), (51^2), (3^21), (32^2), (31^4), (2^21^3), (1^7),$$

and those for $\mathcal{N}(\mathfrak{sp}_6)$ are:

$$(6), (42), (41^2), (3^2), (2^3), (2^21^2), (21^4), (1^6).$$

The Springer correspondence gives a new parametrization of these nilpotent orbits in terms of irreps of the common Weyl group

$$W(B_n) = W(C_n) = \{\pm 1\} \wr S_n.$$

These irreps are labelled by the set \mathcal{Q}_n of **bipartitions** $(\mu; \nu)$ of n .

The Springer correspondence gives a new parametrization of these nilpotent orbits in terms of irreps of the common Weyl group

$$W(B_n) = W(C_n) = \{\pm 1\} \wr S_n.$$

These irreps are labelled by the set \mathcal{Q}_n of **bipartitions** $(\mu; \nu)$ of n .

Theorem (Lusztig, Shoji)

The Springer parametrizations by bipartitions


$$\begin{aligned} SO_{2n+1} \setminus \mathcal{N}(\mathfrak{so}_{2n+1}) &\longleftrightarrow \{(\mu; \nu) \in \mathcal{Q}_n \mid \mu_i \geq \nu_i - 2, \nu_i \geq \mu_{i+1}\} \\ Sp_{2n} \setminus \mathcal{N}(\mathfrak{sp}_{2n}) &\longleftrightarrow \{(\mu; \nu) \in \mathcal{Q}_n \mid \mu_i \geq \nu_i - 1, \nu_i \geq \mu_{i+1} - 1\} \end{aligned}$$

are obtained from the previous by taking 2-quotients of partitions, where the conventions are such that the open orbit corresponds to $((n); \emptyset)$ and the zero orbit corresponds to $(\emptyset; (1^n))$.


Write \mathcal{Q}_n^B and \mathcal{Q}_n^C for these subsets of \mathcal{Q}_n , and $\mathcal{O}_{\mu; \nu}^B$ or $\mathcal{O}_{\mu; \nu}^C$ for the orbit corresponding to $(\mu; \nu) \in \mathcal{Q}_n^B$ or $(\mu; \nu) \in \mathcal{Q}_n^C$ respectively.

Example ($n = 3$, ordering by orbit closure inclusion)

$SO_7 \setminus \mathcal{N}(\mathfrak{so}_7)$



$Sp_6 \setminus \mathcal{N}(\mathfrak{sp}_6)$



Kato's exotic Springer correspondence

The Sp_{2n} -invariant complement of \mathfrak{sp}_{2n} in \mathfrak{gl}_{2n} is

$$S = \{x \in \text{Mat}_{2n} \mid \langle xv, w \rangle = \langle v, xw \rangle, \forall v, w \in \mathbb{C}^{2n}\}.$$

Let $\mathcal{N}(S) = S \cap \mathcal{N}(\mathfrak{gl}_n)$ be its Hilbert nullcone.

Kato's exotic Springer correspondence

The Sp_{2n} -invariant complement of \mathfrak{sp}_{2n} in \mathfrak{gl}_{2n} is

$$S = \{x \in \text{Mat}_{2n} \mid \langle xv, w \rangle = \langle v, xw \rangle, \forall v, w \in \mathbb{C}^{2n}\}.$$

Let $\mathcal{N}(S) = S \cap \mathcal{N}(\mathfrak{gl}_n)$ be its Hilbert nullcone.

Theorem (Kostant, Sekiguchi)

The Sp_{2n} -orbits in $\mathcal{N}(S)$ are classified by Jordan form:

$$Sp_{2n} \backslash \mathcal{N}(S) \longleftrightarrow \left\{ \begin{array}{l} \text{partitions of } 2n \text{ in which every} \\ \text{part occurs with even multiplicity} \end{array} \right\} \longleftrightarrow \mathcal{P}_n.$$

In terms of real groups, $Sp_{2n} \backslash \mathcal{N}(S) \longleftrightarrow GL_n(\mathbb{H}) \backslash \mathcal{N}(\mathfrak{gl}_n(\mathbb{H}))$.

Kato's exotic Springer correspondence

The Sp_{2n} -invariant complement of \mathfrak{sp}_{2n} in \mathfrak{gl}_{2n} is

$$S = \{x \in \text{Mat}_{2n} \mid \langle xv, w \rangle = \langle v, xw \rangle, \forall v, w \in \mathbb{C}^{2n}\}.$$

Let $\mathcal{N}(S) = S \cap \mathcal{N}(\mathfrak{gl}_n)$ be its Hilbert nullcone.

Theorem (Kostant, Sekiguchi)

The Sp_{2n} -orbits in $\mathcal{N}(S)$ are classified by Jordan form:

$$Sp_{2n} \backslash \mathcal{N}(S) \longleftrightarrow \left\{ \begin{array}{l} \text{partitions of } 2n \text{ in which every} \\ \text{part occurs with even multiplicity} \end{array} \right\} \longleftrightarrow \mathcal{P}_n.$$

In terms of real groups, $Sp_{2n} \backslash \mathcal{N}(S) \longleftrightarrow GL_n(\mathbb{H}) \backslash \mathcal{N}(\mathfrak{gl}_n(\mathbb{H}))$.

The natural resolution of $\mathcal{N}(S)$, as in Hesselink's theory, has fibres $\{V_0 \subset V_1 \subset \cdots \subset V_n \mid \dim V_i = 2i, \langle V_i, V_{n-i} \rangle = 0, x(V_i) \subseteq V_{i-1}\}$.

This does give a Springer correspondence, but for S_n not $W(C_n)$.

Syu Kato's **exotic nilpotent cone** of type C_n is

$$\mathfrak{N} := \mathcal{N}(\mathbb{C}^{2n} \oplus \mathcal{S}) = \{(v, x) \mid v \in \mathbb{C}^{2n}, x \in \mathcal{N}(\mathcal{S})\}.$$

The natural resolution ψ of \mathfrak{N} has the following fibre over (v, x) :

$$\{V_0 \subset V_1 \subset \cdots \subset V_{2n} = \mathbb{C}^{2n} \mid \langle V_i, V_{n-i} \rangle = 0, v \in V_n, x(V_i) \subseteq V_i\}.$$

Note that the total space is a bundle over the flag variety Sp_{2n}/B .

Syu Kato's **exotic nilpotent cone** of type C_n is

$$\mathfrak{N} := \mathcal{N}(\mathbb{C}^{2n} \oplus \mathcal{S}) = \{(v, x) \mid v \in \mathbb{C}^{2n}, x \in \mathcal{N}(\mathcal{S})\}.$$

The natural resolution ψ of \mathfrak{N} has the following fibre over (v, x) :

$$\{V_0 \subset V_1 \subset \cdots \subset V_{2n} = \mathbb{C}^{2n} \mid \langle V_i, V_{n-i} \rangle = 0, v \in V_n, x(V_i) \subseteq V_i\}.$$

Note that the total space is a bundle over the flag variety Sp_{2n}/B .

Theorem (Kato, Duke Math. J. 2009)

The endomorphism algebra of $R\psi_\mathbb{C}$ in the derived category on \mathfrak{N} is isomorphic to the group algebra of $W(C_n)$. Moreover, all stabilizers in Sp_{2n} of points in \mathfrak{N} are connected. Hence we have an **exotic Springer correspondence** $Sp_{2n} \setminus \mathfrak{N} \longleftrightarrow \{\text{irreps of } W(C_n)\}$.*

Note that this exotic Springer correspondence is 'cleaner' than the usual Springer correspondence in type C : it is more like type A .

Write $\mathbb{O}_{\mu; \nu}$ for the orbit in \mathfrak{N} corresponding to $(\mu; \nu) \in \mathcal{Q}_n$.

Geometry of the exotic nilpotent cone

Theorem (Achar–H., Advances in Math. 2008)

For $(\rho; \sigma), (\mu; \nu) \in \mathcal{Q}_n$,

$$\mathbb{O}_{\rho; \sigma} \subseteq \overline{\mathbb{O}_{\mu; \nu}} \iff \begin{array}{rcc} \rho_1 & \leq & \mu_1, \\ \rho_1 + \sigma_1 & \leq & \mu_1 + \nu_1, \\ \rho_1 + \sigma_1 + \rho_2 & \leq & \mu_1 + \nu_1 + \mu_2, \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{array}$$

Geometry of the exotic nilpotent cone

Theorem (Achar–H., Advances in Math. 2008)

For $(\rho; \sigma), (\mu; \nu) \in \mathcal{Q}_n$,

$$\mathbb{O}_{\rho; \sigma} \subseteq \overline{\mathbb{O}_{\mu; \nu}} \iff \begin{array}{rcl} \rho_1 & \leq & \mu_1, \\ \rho_1 + \sigma_1 & \leq & \mu_1 + \nu_1, \\ \rho_1 + \sigma_1 + \rho_2 & \leq & \mu_1 + \nu_1 + \mu_2, \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{array}$$

This partial order on \mathcal{Q}_n arose in Shoji's study of 'limit symbols', in which he also defined type- B/C analogues of Kostka polynomials. We conjecture that these give the intersection cohomology:

$$\sum_i \dim IH_{\mathbb{O}_{\rho; \sigma}}^{2i}(\overline{\mathbb{O}_{\mu; \nu}}) q^i \stackrel{?}{=} q^{b(\rho; \sigma) - b(\mu; \nu)} K_{(\mu; \nu), (\rho; \sigma)}(q^{-1}).$$

We proved the special case $\mu = \emptyset$. An obstacle to the general proof is that we do not know a resolution of every orbit closure.

The poset \mathcal{Q}_n contains the posets \mathcal{Q}_n^B and \mathcal{Q}_n^C , corresponding to nilpotent orbits in $\mathcal{N}(\mathfrak{so}_{2n+1})$ and $\mathcal{N}(\mathfrak{sp}_{2n})$ respectively. We can define two partitions of \mathfrak{N} , into **type-B pieces** and **type-C pieces**:

$$\mathfrak{N} = \bigcup_{(\mu;\nu) \in \mathcal{Q}_n^B} \mathbb{T}_{\mu;\nu}^B \quad \text{where} \quad \mathbb{T}_{\mu;\nu}^B = \overline{\mathbb{O}_{\mu;\nu}} \setminus \bigcup_{\substack{(\tau;\nu) \in \mathcal{Q}_n^B \\ (\tau;\nu) < (\mu;\nu)}} \overline{\mathbb{O}_{\tau;\nu}},$$

$$\mathfrak{N} = \bigcup_{(\mu;\nu) \in \mathcal{Q}_n^C} \mathbb{T}_{\mu;\nu}^C \quad \text{where} \quad \mathbb{T}_{\mu;\nu}^C = \overline{\mathbb{O}_{\mu;\nu}} \setminus \bigcup_{\substack{(\tau;\nu) \in \mathcal{Q}_n^C \\ (\tau;\nu) < (\mu;\nu)}} \overline{\mathbb{O}_{\tau;\nu}}.$$

The poset \mathcal{Q}_n contains the posets \mathcal{Q}_n^B and \mathcal{Q}_n^C , corresponding to nilpotent orbits in $\mathcal{N}(\mathfrak{so}_{2n+1})$ and $\mathcal{N}(\mathfrak{sp}_{2n})$ respectively. We can define two partitions of \mathfrak{N} , into **type-B pieces** and **type-C pieces**:

$$\mathfrak{N} = \bigcup_{(\mu;\nu) \in \mathcal{Q}_n^B} \mathbb{T}_{\mu;\nu}^B \quad \text{where} \quad \mathbb{T}_{\mu;\nu}^B = \overline{\mathcal{O}_{\mu;\nu}} \setminus \bigcup_{\substack{(\tau;v) \in \mathcal{Q}_n^B \\ (\tau;v) < (\mu;\nu)}} \overline{\mathcal{O}_{\tau;v}},$$


$$\mathfrak{N} = \bigcup_{(\mu;\nu) \in \mathcal{Q}_n^C} \mathbb{T}_{\mu;\nu}^C \quad \text{where} \quad \mathbb{T}_{\mu;\nu}^C = \overline{\mathcal{O}_{\mu;\nu}} \setminus \bigcup_{\substack{(\tau;v) \in \mathcal{Q}_n^C \\ (\tau;v) < (\mu;\nu)}} \overline{\mathcal{O}_{\tau;v}}.$$

Theorem (Achar–H.–Sommers, arXiv:1001.4283)


1. For every $(\mu; \nu) \in \mathcal{Q}_n^B$, the piece $\mathbb{T}_{\mu;\nu}^B \subset \mathfrak{N}$ is smooth and has the same equivariant cohomology as $\mathcal{O}_{\mu;\nu}^B \subset \mathcal{N}(\mathfrak{so}_{2n+1})$.
2. For every $(\mu; \nu) \in \mathcal{Q}_n^C$, the piece $\mathbb{T}_{\mu;\nu}^C \subset \mathfrak{N}$ is smooth and has the same equivariant cohomology as $\mathcal{O}_{\mu;\nu}^C \subset \mathcal{N}(\mathfrak{sp}_{2n})$.

For “equivariant cohomology”, one can put “number of \mathbb{F}_q -points”.

Example ($n = 3$, type-B pieces)



Example ($n = 3$, type-C pieces)



Why is the nullcone $\mathfrak{N} = \mathcal{N}(\mathbb{C}^{2n} \oplus S) = \mathcal{N}(\mathbb{C}^{2n} \oplus \Lambda^2(\mathbb{C}^{2n}))$ special, and why does it seem just as close to type B as to type C ? A clue is the fact that the nonzero weights of $\mathbb{C}^{2n} \oplus \Lambda^2(\mathbb{C}^{2n})$ form a root system of type B_n , and they each have multiplicity 1.

Why is the nullcone $\mathfrak{N} = \mathcal{N}(\mathbb{C}^{2n} \oplus S) = \mathcal{N}(\mathbb{C}^{2n} \oplus \Lambda^2(\mathbb{C}^{2n}))$ special, and why does it seem just as close to type B as to type C ? A clue is the fact that the nonzero weights of $\mathbb{C}^{2n} \oplus \Lambda^2(\mathbb{C}^{2n})$ form a root system of type B_n , and they each have multiplicity 1.

Proposition

Let \mathbb{F} be an algebraically closed field of characteristic 2. There is a bijective isogeny $SO_{2n+1}(\mathbb{F}) \rightarrow Sp_{2n}(\mathbb{F})$, and short exact sequences:

$$0 \longrightarrow \mathbb{F}^{2n} \longrightarrow \mathfrak{so}_{2n+1}(\mathbb{F}) \longrightarrow \Lambda^2(\mathbb{F}^{2n}) \longrightarrow 0$$

$$0 \longrightarrow \Lambda^2(\mathbb{F}^{2n}) \longrightarrow \mathfrak{sp}_{2n}(\mathbb{F}) \longrightarrow (\mathbb{F}^{2n})^{(1)} \longrightarrow 0$$

So $\mathbb{F}^{2n} \oplus \Lambda^2(\mathbb{F}^{2n})$ is an 'exotic adjoint representation', obtained by degenerating $\mathfrak{so}_{2n+1}(\mathbb{F})$ or $\mathfrak{sp}_{2n}(\mathbb{F})$. Similarly, $\mathfrak{N}(\mathbb{F})$ can be viewed as a degeneration of $\mathcal{N}(\mathfrak{so}_{2n+1}(\mathbb{F}))$ or of $\mathcal{N}(\mathfrak{sp}_{2n}(\mathbb{F}))$. This implies the desired equalities of the numbers of \mathbb{F}_q -points when $q = 2^s$.

To prove smoothness of the type- B and type- C pieces of \mathfrak{N} in any characteristic, we gave resolutions of $\overline{\mathbb{O}}_{\mu;\nu}$ for $(\mu;\nu) \in \mathcal{Q}_n^B \cup \mathcal{Q}_n^C$. These were inspired by the Jacobson–Morozov resolutions of $\overline{\mathcal{O}}_{\mu;\nu}^B$ and $\overline{\mathcal{O}}_{\mu;\nu}^C$, and are in the spirit of Hesselink’s general theory.

To prove smoothness of the type- B and type- C pieces of \mathfrak{N} in any characteristic, we gave resolutions of $\overline{\mathcal{O}}_{\mu;\nu}$ for $(\mu; \nu) \in \mathcal{Q}_n^B \cup \mathcal{Q}_n^C$. These were inspired by the Jacobson–Morozov resolutions of $\overline{\mathcal{O}}_{\mu;\nu}^B$ and $\overline{\mathcal{O}}_{\mu;\nu}^C$, and are in the spirit of Hesselink’s general theory.

Let $\mathcal{Q}_n^\circ = \mathcal{Q}_n^B \cap \mathcal{Q}_n^C$ be the set labelling Lusztig’s special irreps. All cones have a partition into **special pieces** indexed by $(\mu; \nu) \in \mathcal{Q}_n^\circ$:

$$\mathcal{N}(\mathfrak{so}_{2n+1}) = \bigcup \mathcal{S}_{\mu;\nu}^B, \quad \mathcal{N}(\mathfrak{sp}_{2n}) = \bigcup \mathcal{S}_{\mu;\nu}^C, \quad \mathfrak{N} = \bigcup \mathcal{S}_{\mu;\nu}.$$

It follows immediately from our result that $\mathcal{S}_{\mu;\nu}^B$, $\mathcal{S}_{\mu;\nu}^C$, and $\mathcal{S}_{\mu;\nu}$ have the same equivariant cohomology (or number of \mathbb{F}_q -points). Lusztig had already proved this for $\mathcal{S}_{\mu;\nu}^B$ and $\mathcal{S}_{\mu;\nu}^C$.

To prove smoothness of the type- B and type- C pieces of \mathfrak{N} in any characteristic, we gave resolutions of $\overline{\mathcal{O}}_{\mu;\nu}$ for $(\mu;\nu) \in \mathcal{Q}_n^B \cup \mathcal{Q}_n^C$. These were inspired by the Jacobson–Morozov resolutions of $\overline{\mathcal{O}}_{\mu;\nu}^B$ and $\overline{\mathcal{O}}_{\mu;\nu}^C$, and are in the spirit of Hesselink’s general theory.

Let $\mathcal{Q}_n^\circ = \mathcal{Q}_n^B \cap \mathcal{Q}_n^C$ be the set labelling Lusztig’s special irreps. All cones have a partition into **special pieces** indexed by $(\mu;\nu) \in \mathcal{Q}_n^\circ$:

$$\mathcal{N}(\mathfrak{so}_{2n+1}) = \bigcup \mathcal{S}_{\mu;\nu}^B, \quad \mathcal{N}(\mathfrak{sp}_{2n}) = \bigcup \mathcal{S}_{\mu;\nu}^C, \quad \mathfrak{N} = \bigcup \mathcal{S}_{\mu;\nu}.$$

It follows immediately from our result that $\mathcal{S}_{\mu;\nu}^B$, $\mathcal{S}_{\mu;\nu}^C$, and $\mathcal{S}_{\mu;\nu}$ have the same equivariant cohomology (or number of \mathbb{F}_q -points). Lusztig had already proved this for $\mathcal{S}_{\mu;\nu}^B$ and $\mathcal{S}_{\mu;\nu}^C$.

Theorem (Kraft–Procesi)

In characteristic $\neq 2$, $\mathcal{S}_{\mu;\nu}^B$ and $\mathcal{S}_{\mu;\nu}^C$ are rationally smooth.

So it is natural to conjecture that $\mathcal{S}_{\mu;\nu}$ is rationally smooth (or possibly smooth?) in any characteristic.