

Stability conditions on Dia

D 0 Every finite partially ordered set is in Dia.

D 1 Dia is stable under finite sums and fiber products.

D 2d If I is in Dia and i an object of I , then $i \setminus I$ is in Dia.

D 2g If I is in Dia and i an object of I , then I/i is in Dia.

D 3 If I is in Dia, then I^0 is in Dia.

D 4d If $u: I \rightarrow J$ is a filtered category on J (in Grothendieck's sense) and if J and the fibers of u are in Dia, then I is in Dia.

D 4g If $u: I \rightarrow J$ is a filtered category on J (in Grothendieck's sense) and if J and the fibers of u are in Dia, then I is in Dia.

(D 3) \Rightarrow ((D 2d) \Leftrightarrow (D 2g)) and (D 4d) \Leftrightarrow (D 4g)

(D 4g) \Rightarrow (D 2g) and (D 4d) \Rightarrow D 2d if discrete categories (small) are in Dia.

(D 1) and (D 2d) \Rightarrow If $I \rightarrow J$ is in Dia and j an object of J , then $j \setminus I$ is in Dia.

(D 1) and (D 2g) \Rightarrow If $I \rightarrow J$ is in Dia and j an object of J , then I/j is in Dia.

Axioms of Derivators

Let $\mathcal{D}ia$ a full 2-subcategory of $\mathcal{C}at$ satisfying all stability conditions needed and

$$\mathbb{D}: \mathcal{D}ia^{\circ} \longrightarrow \mathcal{C}AT$$

a 2-functor (a prederivator). The prederivator \mathbb{D} is called a derivator if the following five axioms are satisfied

Der 1] i) The category $\mathbb{D}(\emptyset)$ is equivalent to the point category \mathbf{e} .

ii) Let I, J in $\mathcal{D}ia$ and

$$\begin{array}{ccc} I & \xrightarrow{i} & I \amalg J \\ J & \xrightarrow{j} & I \amalg J \end{array}$$

the canonical maps. Then

$$\mathbb{D}(I \amalg J) \xrightarrow{[i^*, j^*]} \mathbb{D}(I) \times \mathbb{D}(J)$$

is an equivalence of categories

Der 2] For every I in $\mathcal{D}ia$ the family of functors

$$i^*: \mathbb{D}(I) \longrightarrow \mathbb{D}(e), \quad i \in \mathcal{O}b I,$$

where $i: e \rightarrow I$ denotes also the functor defined by the object i , is conservative

Der 3a] For every $u: I \rightarrow J$ in $\mathcal{D}ia$ the functor $u^*: \mathbb{D}(J) \rightarrow \mathbb{D}(I)$ has a left adjoint $u_!: \mathbb{D}(I) \rightarrow \mathbb{D}(J)$

Der 3b] For every $u: I \rightarrow J$ in $\mathcal{D}ia$ the functor $u^*: \mathbb{D}(J) \rightarrow \mathbb{D}(I)$ has a right adjoint $u_*: \mathbb{D}(I) \rightarrow \mathbb{D}(J)$

Der 4a For every $u: I \rightarrow J$ in $\mathcal{D}ia$ and every object j of \mathcal{J} the canonical map

$$j^* u_! \leftarrow p_! k^*$$

induced by the "2-square"

$$\begin{array}{ccc} j \backslash I & \xrightarrow{k} & I \\ p \downarrow & \nearrow & \downarrow u \\ e & \xrightarrow{j} & J \end{array}$$

is an isomorphism

Der 4a For every $u: I \rightarrow J$ in $\mathcal{D}ia$ and every object j of \mathcal{J} the canonical map

$$j^* u_* \rightarrow q_* l^*$$

induced by the "2-square"

$$\begin{array}{ccc} I/j & \xrightarrow{l} & I \\ q \downarrow & \searrow & \downarrow u \\ e & \xrightarrow{j} & J \end{array}$$

is an isomorphism

Der 5 For every I in $\mathcal{D}ia$ the canonical functor

$$\mathcal{D}(I \times \{0 \rightarrow 1\}) \rightarrow \underline{\text{Hom}}(\{0 \rightarrow 1\}^{\circ}, \mathcal{D}(I))$$

is full and essentially surjective

Additional axioms for triangulated derivators

Der 6 If $u: I \rightarrow J$ is an open (resp. a closed) immersion in $\mathcal{D}ia$, then the functor $u_!$ (resp. u_*) has a left (resp. right) adjoint $u^?$ (resp. $u^!$)

Der 7 An object of $\mathcal{D}(I) = \mathcal{D}\left(\begin{array}{ccc} (0,0) & \xrightarrow{k} & (0,1) \\ \uparrow & & \uparrow \\ (1,0) & \xrightarrow{l} & (1,1) \end{array}\right)$ is cocartesian if and only if it is cartesian.

Results on \mathbb{D} -equivalences, \mathbb{D} -aspheric maps, etc.

A) $u: I \rightarrow J$ in $\mathcal{D}\text{ca}$. Equivalent conditions

a) u \mathbb{D} -equivalence;

b) $p_I!: p_I^* \rightarrow p_J!: p_J^*$ isomorphism;

c) $p_J^* \rightarrow p_I^* p_I^*$ isomorphism;

B) $u: I \rightarrow J$ in $\mathcal{D}\text{ca}$. Equivalent conditions

a) u \mathbb{D} -aspheric

b) $p_J^* \rightarrow u_* p_I^*$ isomorphism;

c) $(p_I)!: u^* \rightarrow (p_J)!$ isomorphism;

d) $\forall j \in \text{Ob } J, I/j \rightarrow J/j$ is a \mathbb{D} -equivalence.

C) $u: I \rightarrow J$ in $\mathcal{D}\text{ca}$. Equivalent conditions

a) u \mathbb{D} -coaspheric

b) $u!: p_I^* \rightarrow p_J^*$ isomorphism;

c) $(p_J)_* \rightarrow (p_I)_* u^*$ isomorphism;

d) $\forall j \in \text{Ob } J, j \setminus I \rightarrow j \setminus J$ is a \mathbb{D} -equivalence.

D)
$$\begin{array}{ccc} I & \xrightarrow{u} & J \\ \searrow v & & \swarrow w \\ & K & \end{array}$$
 commutative triangle in $\mathcal{D}\text{ca}$

Equivalent conditions:

a) u is a \mathbb{D} -equivalence locally on K ;

b) $w_* p_J^* \rightarrow v_* p_I^*$ isomorphism;

c) $(p_I)!: v^* \rightarrow (p_J)!: w^*$ isomorphism;

d) $\forall k \in \text{Ob } K, I/k \rightarrow J/k$ is a \mathbb{D} -equivalence.

E)
$$\begin{array}{ccc} I & \xrightarrow{u} & J \\ v \searrow & & \swarrow w \\ & K & \end{array}$$
 commutative triangle in $\mathcal{D}(\mathcal{A})$
Equivalent conditions:

- u is a \mathbb{D} -equivalence colocally on K ;
- $\forall ! p_I^* \rightarrow w ! p_J^*$ isomorphism;
- $(p_J)_* w^* \rightarrow (p_I)_* v^*$ isomorphism;
- $\forall R \in \text{Ob } K, R \setminus I \rightarrow R \setminus J$ is a \mathbb{D} -equivalence.

F) I in $\mathcal{D}(\mathcal{A})$. Equivalent conditions

- I \mathbb{D} -aspheric \Rightarrow
- p_I \mathbb{D} -equivalence;
- p_I \mathbb{D} -aspheric;
- p_I \mathbb{D} -coaspheric;
- $(p_I)_! p_I^* \rightarrow 1_{\mathbb{D}(\mathcal{A})}$ isomorphism;
- $1_{\mathbb{D}(\mathcal{A})} \rightarrow (p_I)_* p_I^*$ isomorphism;
- p_I^* fully faithful.

G) Duality:

- $u: I \rightarrow J$ \mathbb{D} -equivalence $\Leftrightarrow u^\circ: I^\circ \rightarrow J^\circ$ \mathbb{D} -equivalence;
- $u: I \rightarrow J$ \mathbb{D} -aspheric $\Leftrightarrow u^\circ: I^\circ \rightarrow J^\circ$ \mathbb{D} -coaspheric;

c)
$$\begin{array}{ccc} I & \xrightarrow{u} & J \\ \searrow & & \swarrow \\ & K & \end{array}$$
 commutative triangle

u \mathbb{D} -equivalence locally on $K \Leftrightarrow$

$\Leftrightarrow u^\circ$ \mathbb{D} -equivalence colocally on K

d) I \mathbb{D} -aspheric $\Leftrightarrow I^\circ$ \mathbb{D} -aspheric

H) $\mathcal{D} = \begin{array}{ccc} I' & \xrightarrow{v} & I \\ u \downarrow & \swarrow \alpha & \downarrow u \\ J' & \xrightarrow{w} & J \end{array}$ 2-square in $\mathcal{D}ia$

$$\mathcal{D} = \begin{array}{ccc} I' & \xrightarrow{v} & I \\ u \downarrow & \swarrow \alpha & \downarrow u \\ J' & \xrightarrow{w} & J \end{array}$$

2-square in $\mathcal{D}ia$

Equivalent conditions:

- \mathcal{D} is \mathcal{D} -exact;
- $c_{\mathcal{D}}: v_! u^* \rightarrow u^* w_!$ isomorphism;
- $c'_{\mathcal{D}}: w^* u_* \rightarrow u'_* v^*$ isomorphism;
- $\forall i \in \text{ob } I, i \setminus I' \rightarrow u(i) \setminus J'$ is a \mathcal{D} -equivalence locally on J' ;
- $\forall j' \in \text{ob } J', I'/j' \rightarrow I/w(j')$ is a \mathcal{D} -equivalence colocally on I .

I) $u: I \rightarrow J$ in $\mathcal{D}ia$. Equivalent conditions:

- u is \mathcal{D} -proper;
- for every diagram of cartesian squares of the form

$$\begin{array}{ccccc} I'' & \longrightarrow & I' & \longrightarrow & I \\ \downarrow & & \downarrow & & \downarrow u \\ J'' & \longrightarrow & J' & \longrightarrow & J \end{array}$$

the left square is \mathcal{D} -exact;

- for every diagram of cartesian squares of the form

$$\begin{array}{ccccc} I'' & \xrightarrow{v} & I' & \longrightarrow & I \\ \downarrow & & \downarrow & & \downarrow u \\ J'' & \xrightarrow{w} & J' & \longrightarrow & J \end{array}$$

if w is \mathcal{D} -coyberic then v is \mathcal{D} -coyberic.

- $\forall j \in \text{ob } J, I_j \rightarrow I/j$ is \mathcal{D} -coyberic

- $\forall i \in \text{ob } I, i \setminus I \rightarrow u(i) \setminus J$ has

\mathcal{D} -coyberic fibers

- $\forall i \in \text{ob } I, i \setminus I \rightarrow u(i) \setminus J$ is a \mathcal{D} -equivalence and remains a \mathcal{D} -equivalence after any base change

J) $u: I \rightarrow J$ in $\mathcal{D}ia$. Equivalent conditions:

a) u is \mathbb{D} -smooth

b) for every diagram of cartesian squares of the form

$$\begin{array}{ccc} I'' & \longrightarrow & J'' \\ \downarrow & & \downarrow \\ I' & \longrightarrow & J' \\ \downarrow & & \downarrow \\ I & \xrightarrow{u} & J \end{array} \quad \text{the upper square is } \mathbb{D}\text{-exact};$$

c) for every diagram of cartesian squares of the form

$$\begin{array}{ccccc} I'' & \xrightarrow{v} & I' & \longrightarrow & I \\ \downarrow & & \downarrow & & \downarrow u \\ J'' & \xrightarrow{w} & J' & \longrightarrow & J \end{array} \quad \begin{array}{l} \text{if } w \text{ is } \mathbb{D}\text{-acyclic} \\ \text{then } v \text{ is } \mathbb{D}\text{-acyclic}; \end{array}$$

d) $\forall j \in \text{ob } J, \quad I_j \rightarrow j \setminus I$ is \mathbb{D} -acyclic;

e) $\forall i \in \text{ob } I, \quad I/i \rightarrow J/u(i)$ has \mathbb{D} -acyclic fibers;

f) $\forall i \in \text{ob } I, \quad I/i \rightarrow J/u(i)$ is a \mathbb{D} -equivalence and remains a \mathbb{D} -equivalence after any base change.

K) Duality: $u: I \rightarrow J$ in $\mathcal{D}ia$

$$u: I \rightarrow J \text{ } \mathbb{D}\text{-proper} \iff u^\circ: I^\circ \rightarrow J^\circ \text{ } \mathbb{D}\text{-smooth.}$$